

COMPRESSION TECHNIQUES FOR BOUNDARY INTEGRAL EQUATIONS—ASYMPTOTICALLY OPTIMAL COMPLEXITY ESTIMATES*

WOLFGANG DAHMEN[†], HELMUT HARBRECHT[‡], AND REINHOLD SCHNEIDER[‡]

Abstract. Matrix compression techniques in the context of wavelet Galerkin schemes for boundary integral equations are developed and analyzed that exhibit optimal complexity in the following sense. The fully discrete scheme produces approximate solutions within discretization error accuracy offered by the underlying Galerkin method at a computational expense that is proven to stay proportional to the number of unknowns. Key issues are the second compression, which reduces the near field complexity significantly, and an additional a posteriori compression. The latter is based on a general result concerning an optimal work balance that applies, in particular, to the quadrature used to compute the compressed stiffness matrix with sufficient accuracy in linear time.

Key words. wavelets, norm equivalences, multilevel preconditioning, first and second compression, a posteriori compression, asymptotic complexity estimates

AMS subject classifications. 47A20, 65F10, 65F50, 65N38, 65R20

DOI. 10.1137/S0036142903428852

1. Introduction. Many mathematical models concerning, e.g., field calculations, flow simulation, elasticity, or visualization are based on operator equations with *global operators*, especially *boundary integral operators*. Discretizing such problems will then lead in general to possibly very large linear systems with *densely populated* matrices. Moreover, the involved operator may have an order different from zero, which means that it acts on different length scales in a different way. This is well known to entail the linear systems to become more and more ill-conditioned when the level of resolution increases. Both features pose serious obstructions to the efficient numerical treatment of such problems to an extent that desirable realistic simulations are still beyond the current computing capacities.

This fact has stimulated enormous efforts to overcome these obstructions. The resulting significant progress made over the past 10 or 15 years manifests itself in several different approaches such as panel clustering (PC) [18], multipole expansions (FMM) [16], and wavelet compression (WC) [2]. Each of these methodologies has its recognized advantages and drawbacks whose balance may depend on the problem at hand. The first two (PC, FMM) are quite similar in spirit and exploit perhaps in the best way the (typical) smoothness of the potential kernel in the space rather than the integral kernel on the boundary manifold. As a consequence, they are fairly robust with regard to the shape and complexity of the boundary manifold. This is also the case for further developments like hierarchical matrices (\mathcal{H} -matrices) [17] and adaptive cross approximation (ACA) [1] based on a low rank approximation in the far

*Received by the editors May 28, 2003; accepted for publication (in revised form) March 29, 2005; published electronically January 6, 2006. This work was supported in part by the European Community's Human Potential Programme under contract HPRN-CT-202-00286 (BREAKING COMPLEXITY), and the SFB 393 "Numerical Simulation on Massive Parallel Computers" funded by the Deutsche Forschungsgemeinschaft.

<http://www.siam.org/journals/sinum/43-6/42885.html>

[†]RTWH Aachen, Institut für Geometrie und Praktische Mathematik, Aachen 52056, Germany (dahmen@igpm.rwth-aachen.de).

[‡]Christian-Albrechts-Universität zu Kiel, Institut für Informatik und Praktische Mathematik, Olshausenstr. 40, Kiel 24098, Germany (hh@numerik.uni-kiel.de, rs@numerik.uni-kiel.de).

field; see also [30]. Common experience seems to indicate that the third option (WC) depends in a more sensitive way on the underlying geometry, and its performance may suffer from strong domain anisotropies. On the other hand, WC allows one in a natural way to incorporate preconditioning techniques, which very much supports the fast solution of the resulting sparsified systems. Moreover, recent developments suggest a natural combination with adaptive discretizations to keep from the start, for a given target accuracy, the size of the linear systems as small as possible. Perhaps the main difference between PC, FMM on the one hand and WC on the other is that the former are essentially *agglomeration* techniques, while WC is more apt for refining a given coarse discretization. Since these methodologies are, in that sense, somewhat complementary in spirit, it is in our opinion less appropriate to contrapose them, but one should rather try to extract the best from each option.

As indicated before, a preference for any of the above-mentioned solution strategies will, in general, depend on the concrete application at hand. The objective of this paper is therefore to provide a complete analysis of the wavelet approach (WC) from the following perspectives. Recall that WC has been essentially initiated by the pioneering paper [2], where it was observed that certain operators have an almost sparse representation in wavelet coordinates. Discarding all entries below a certain threshold in a given principal section of the wavelet representation will then give rise to a sparse matrix that can be further processed by efficient linear algebra tools. This idea has since then initiated many subsequent studies. The diversity as well as the partly deceiving nature of the by now existing rich body of literature is one reason for us to take up this subject here again. Our attempt to provide a unified analysis is certainly based, to some extent, on previously used techniques. However, on the one hand we have extended these concepts essentially by several new analytical tools, to be detailed later below. On the other hand, the numerical implementation has been brought now to a state that makes the method applicable to practically relevant problems. The main objective of this paper is to present these new theoretical and practical developments centering on the following issues.

When dealing with large-scale problems, a sharp *asymptotic analysis* of the complexity is in our opinion ultimately essential for assessing its potential. It is important to clarify the meaning of “complexity” in this context. It is always understood as the *work/accuracy* rate of the method under consideration when the level of resolution increases, i.e., the overall accuracy of the computed approximate solution is to be tied to the computational work required to obtain it. There is no point in increasing the number of degrees of freedom, i.e., the size of the linear systems, without improving the accuracy of the resulting solutions. On the other hand, since one is ultimately interested in the “exact solution of the infinite-dimensional problem” it makes no sense to determine the solution to the discrete finite-dimensional problem with much more accuracy than that offered by the discretization. Thus, a reasonable target accuracy for the solutions of the discrete systems is *discretization error accuracy* which will guide all our subsequent considerations. A method will therefore be said to exhibit *asymptotically optimal complexity* if the discrete solution can be computed, for any discretization level, within discretization error accuracy at a computational expense that stays *proportional* to the number N of unknowns, i.e., when not even logarithmic factors are permitted. Obviously, computational optimality refers here to the given discretization framework based on the chosen hierarchy of trial spaces.

In the present context, this means that the solutions of the compressed systems should exhibit the same asymptotic convergence rates as the solutions to unperturbed discretizations. In connection with WC, this in turn means that any threshold param-

eters and truncation strategies have to be adapted to the current number N of degrees of freedom. Such an asymptotic analysis is missing in [2] and in many subsequent investigations. The program carried out in [11, 12, 24, 25, 27] aimed at determining exactly such work/accuracy rates for various types of boundary integral equations. Roughly speaking, it could be shown that discretization error accuracy can be realized for appropriately chosen wavelet bases at a computational work that stays bounded by $C N(\log N)^a$ for some constants C, a independent of the size N of the linear systems. Moreover, in [27] it was shown for the first time that, by incorporating a second compression, an overall optimal compression strategy can be devised that even avoids additional logarithmic factors, while the complexity estimates for a corresponding adaptive quadrature scheme were confined to collocation methods.

The purpose of the present paper can now be summarized as follows.

We present a complete, in several respects new and improved analysis of wavelet compression schemes for boundary integral equations based on Galerkin discretizations that exhibit overall asymptotically optimal complexity. This means that discretization error accuracy is obtained at a computational expense that stays proportional to the size N of the arising linear systems, uniformly in N . In contrast to the earlier treatments of the authors it is based on bilinear forms and Strang's lemma; see also [25].

The analysis significantly simplifies previous studies including the effect of the second compression. In fact, the analysis of the second compression is completely new, resulting in slightly different conclusions; see section 6. The complete work balance synchronizing quadrature and compression accuracy is based on new *balance estimates* given in section 11; see Theorem 11.1. In particular, it reveals the right work balance for the compression and the quadrature needed to compute the compressed matrices with sufficient accuracy, so as to realize asymptotically optimal computational complexity of the fully discretized scheme. Specifically, the computational work for computing and assembling the compressed stiffness matrix remains proportional to the number N of degrees of freedom.

This also lays the foundation for an additional new *a posteriori compression* whose analysis is again based on the above-mentioned balance estimate. This improves the quantitative performance of the scheme significantly as shown in [20, 22].

Our analysis concerns what is called the *standard* wavelet representation. A preference for using the so-called *nonstandard* form is frequently reported in the literature. The reason is that the entries in this latter form only involve scaling functions and wavelets on the *same* level. This indeed simplifies assembling the matrices and offers essential advantages when dealing with shift-invariant problems. However, aside from the problem of preconditioning in connection with operators of nonzero order, to our knowledge it has so far *not* been shown that, for a fixed order of vanishing moments, optimal computational complexity in the above sense can be obtained with the nonstandard form. In fact, for regular solutions approximate solutions with prescribed accuracy can be obtained at (asymptotically) a lower computational cost with the aid of the standard form when compared with the nonstandard form. This is backed by theory and confirmed by numerical experience; see [20, 22].

As mentioned before, it is important to employ the "right" wavelet bases. This question has been discussed extensively in previous work [3, 6, 13, 14]. The theory tells us that, depending on the order of the operator, a proper relation between the approximation order of the underlying multiresolution spaces and the order of vanishing moments matters, which often rules out orthonormal wavelets. Given the validity

of this relation it is important to keep supports as small as possible; see Remark 7.1 in section 7. Moreover, our present analysis of the second compression refers exclusively to biorthogonal spline wavelets whose singular supports are well defined; see [21] for examples and graphical illustrations.

We shall frequently write $a \lesssim b$ to express that a is bounded by a constant multiple of b , uniformly with respect to all parameters on which a and b may depend. Then $a \sim b$ means $a \lesssim b$ and $b \lesssim a$.

2. Problem formulation and preliminaries. We consider boundary integral equations on a closed boundary surface Γ of an $(n + 1)$ -dimensional domain $\Omega \subset \mathbb{R}^{n+1}$

$$(2.1) \quad Au(\hat{x}) = \int_{\Gamma} k(\hat{x}, \hat{y})u(\hat{y})d\Gamma_{\hat{y}} = f(\hat{x}), \quad \hat{x} \in \Gamma,$$

where the boundary integral operator is assumed to be an operator of order $2q$, that is, $A : H^q(\Gamma) \rightarrow H^{-q}(\Gamma)$. The kernel functions under consideration are supposed to be smooth as functions in the variables \hat{x}, \hat{y} , apart from the diagonal $\{(\hat{x}, \hat{y}) \in \Gamma \times \Gamma : \hat{x} = \hat{y}\}$ and may have a singularity on the diagonal. Such kernel functions arise, for instance, by applying a boundary integral formulation to a second-order elliptic problem. In general, they decay like a negative power of the distance of the arguments which depends on the spatial dimension n and the order $2q$ of the operator.

Throughout the remainder of this paper we shall assume that the boundary manifold Γ is given as a parametric surface consisting of smooth patches. More precisely, let $\square := [0, 1]^n$ denote the unit n cube. The manifold $\Gamma \subset \mathbb{R}^{n+1}$ is partitioned into a finite number of *patches*,

$$(2.2) \quad \Gamma = \bigcup_{i=1}^M \Gamma_i, \quad \Gamma_i = \gamma_i(\square), \quad i = 1, 2, \dots, M,$$

where each $\gamma_i : \square \rightarrow \Gamma_i$ defines a diffeomorphism of \square onto Γ_i . We also assume that there exist smooth extensions $\tilde{\Gamma}_i \subset \subset \tilde{\Gamma}$ and $\tilde{\gamma}_i : \tilde{\square} := [-1, 2]^n \rightarrow \tilde{\Gamma}_i$. The intersection $\Gamma_i \cap \Gamma_{i'}, i \neq i'$, of the patches Γ_i and $\Gamma_{i'}$ is supposed to be either \emptyset or a lower-dimensional face.

A mesh of level j on Γ is induced by dyadic subdivisions of depth j of the unit cube into 2^{nj} cubes $C_{j,k} \subseteq \square$, where $k = (k_1, \dots, k_n)$ with $0 \leq k_m < 2^j$. This generates $2^{nj}M$ *elements* (or elementary domains) $\Gamma_{i,j,k} := \gamma_i(C_{j,k}) \subseteq \Gamma_i, i = 1, \dots, M$.

In order to ensure that the collection of elements $\{\Gamma_{i,j,k}\}$ on the level j forms a regular mesh on Γ , the parametric representation is subjected to the following *matching condition*: for all $\hat{x} \in \Gamma_i \cap \Gamma_{i'}$ there exists a bijective, affine mapping $\Xi : \square \rightarrow \square$ such that $\gamma_i(x) = (\gamma_{i'} \circ \Xi)(x) = \hat{x}$ for $x = (x_1, \dots, x_n) \in \square$ with $\gamma_i(x) = \hat{x}$.

The first fundamental tensor of differential geometry is given by the matrix $K_i(x) \in \mathbb{R}^{n \times n}$ defined by

$$(2.3) \quad K_i(x) := \left[\left(\frac{\partial \gamma_i(x)}{\partial x_j}, \frac{\partial \gamma_i(x)}{\partial x_{j'}} \right)_{l^2(\mathbb{R}^{n+1})} \right]_{j,j'=1}^n.$$

Since γ_i is supposed to be a diffeomorphism, the matrix $K_i(x)$ is symmetric and positive definite. The canonical inner product in $L^2(\Gamma)$ is then given by

$$(2.4) \quad \langle u, v \rangle = \int_{\Gamma} u(x)v(x)d\Gamma_x = \sum_{i=1}^M \int_{\square} u(\gamma_i(x))v(\gamma_i(x))\sqrt{\det K_i(x)}dx.$$

The corresponding Sobolev spaces are denoted by $H^s(\Gamma)$, endowed with the norms $\|\cdot\|_s$, where for $s < 0$ it is understood that $H^s(\Gamma) = (H^{-s}(\Gamma))'$. Of course, depending on the global smoothness of the surface, the range of permitted $s \in \mathbb{R}$ is limited to $s \in (-s_\Gamma, s_\Gamma)$. In the case of general Lipschitz domains we have at least $s_\Gamma = 1$ since for all $0 \leq s \leq 1$ the spaces $H^s(\Gamma)$ consist of traces of functions $\in H^{s+1/2}(\Omega)$; cf. [7].

We can now specify the kernel functions. To this end, we denote by $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\beta = (\beta_1, \dots, \beta_n)$ multi-indices of dimension n and define $|\alpha| := \alpha_1 + \dots + \alpha_n$. Recall that \hat{x} and \hat{y} are points on the surface, i.e., $\hat{x} := \gamma_i(x)$ and $\hat{y} := \gamma_{i'}(y)$ for some $1 \leq i, i' \leq M$.

DEFINITION 2.1. *A kernel $k(\hat{x}, \hat{y})$ is called standard kernel of order $2q$ if the partial derivatives of the transported kernel function*

$$(2.5) \quad \tilde{k}(x, y) := k(\gamma_i(x), \gamma_{i'}(y)) \sqrt{\det K_i(x)} \sqrt{\det K_{i'}(y)}$$

are bounded by

$$(2.6) \quad |\partial_x^\alpha \partial_y^\beta \tilde{k}(x, y)| \leq c_{\alpha, \beta} \text{dist}(\hat{x}, \hat{y})^{-(n+2q+|\alpha|+|\beta|)},$$

provided that $n + 2q + |\alpha| + |\beta| > 0$.

We emphasize that this definition requires patchwise smoothness but *not* global smoothness of the geometry. The surface itself needs to be only Lipschitz. Generally, under this assumption, the kernel of a boundary integral operator A of order $2q$ is a standard kernel of order $2q$. Hence, we may assume this property in the following. We shall encounter further specifications below in connection with discretizations.

3. Galerkin scheme. We shall be concerned with the Galerkin method with respect to a hierarchy of conforming trial spaces $V_J \subset V_{J+1} \subset H^q(\Gamma)$: find $u_J \in V_J$ solving the variational problem

$$(3.1) \quad \langle Au_J, v_J \rangle = \langle f, v_J \rangle \quad \text{for all } v_J \in V_J.$$

Here the index J reflects a meshwidth of the order 2^{-J} . Moreover, we say that the trial spaces have (*approximation*) order $d \in \mathbb{N}$ and (*regularity*) $\gamma > 0$ if

$$(3.2) \quad \begin{aligned} \gamma &= \sup\{s \in \mathbb{R} : V_J \subset H^s(\Gamma)\}, \\ d &= \sup\{s \in \mathbb{R} : \inf_{v_J \in V_J} \|v - v_J\|_0 \lesssim 2^{-Js} \|v\|_s \text{ for all } v \in H^s(\Gamma)\}. \end{aligned}$$

Thus conformity requires, of course, that $\gamma > \max\{0, q\}$.

In order to ensure that (3.1) is well posed we shall make the following assumptions on the operator A throughout the remainder of the paper.

Assumptions:

1. A is strongly elliptic, i.e., there exists a symmetric compact operator $C : H^q(\Gamma) \rightarrow H^{-q}(\Gamma)$ such that $\langle (A + A^* + C)u, u \rangle \gtrsim \|u\|_q^2$.
2. The operator $A : H^q(\Gamma) \rightarrow H^{-q}(\Gamma)$ is injective, i.e., $\text{Ker } A = \{0\}$.

Remark. 1. Most boundary integral equations of the first kind, resulting from a direct approach, are known to be strongly elliptic, even if Γ is supposed to be the boundary of a Lipschitz domain [7]. In particular, this is the case for boundary integral equations of the first kind for the Laplacian, the system of Navier–Lamé equations, and the Stokes system. For integral equations of the second kind the condition is obvious if the double layer potential operator is compact, or in the case of smooth boundaries, since the principal symbol satisfies a Gårding inequality.

2. For several boundary integral operators like the single layer operator of the Stokes system, for operators associated with Neumann problems or multiply connected domains, the second assumption is not valid. But in these cases the kernel of the operator A is finite-dimensional and known a priori. The kernels can be factored out, i.e., $A : H^q(\Gamma)/\text{Ker}(A) \rightarrow (H^q(\Gamma)/\text{Ker}(A))'$. A standard approach uses constrained conditions and Lagrange multipliers. With a minor modification our method can be applied also to these cases. Therefore, the second assumption is only for the sake of simplicity and not a restriction of generality.

LEMMA 3.1 (see, e.g., [31]). *Under the above assumptions the Galerkin discretization is stable, i.e.,*

$$(3.3) \quad \langle (A + A^*)v_J, v_J \rangle \gtrsim \|v_J\|_q^2, \quad v_J \in V_J,$$

for J sufficiently large, and

$$(3.4) \quad |\langle Av_J, w_J \rangle| \lesssim \|v_J\|_q \|w_J\|_q, \quad v_J, w_J \in V_J.$$

Furthermore, let u, u_J denote the solution of the original equation $Au = f$, respectively, of (3.1). Then one has

$$(3.5) \quad \|u - u_J\|_t \lesssim 2^{J(t-t')} \|u\|_{t'}$$

provided that $2q - d \leq t < \gamma$, $t \leq t'$, $q \leq t' \leq d$ and Γ is sufficiently regular.

Note that the best possible convergence rate is given by

$$(3.6) \quad \|u - u_J\|_{2q-d} \lesssim 2^{-2J(d-q)} \|u\|_d$$

provided that $u \in H^d(\Gamma)$, which is only possible when Γ is sufficiently regular. Since this case gives rise to the highest convergence rate, it will be seen later to impose the most stringent demands on the matrix compression.

4. Wavelets and multiresolution analysis. The nested trial spaces $V_j \subset V_{j+1}$ that we shall employ in (3.1) are spanned by so-called *single-scale bases* $\Phi_j = \{\phi_{j,k} : k \in \Delta_j\}$, where Δ_j denote suitable index sets of cardinality $\dim V_j$. The elements of Φ_j are normalized in $L^2(\Gamma)$ and their compact supports scale like $\text{diam supp } \phi_{j,k} \sim 2^{-j}$. Associated with these collections are always *dual bases* $\tilde{\Phi}_j = \{\tilde{\phi}_{j,k} : k \in \Delta_j\}$, i.e., one has $\langle \phi_{j,k}, \tilde{\phi}_{j,k'} \rangle = \delta_{k,k'}$, $k, k' \in \Delta_j$. For the current type of boundary surfaces Γ the $\Phi_j, \tilde{\Phi}_j$ are generated by constructing first dual pairs of single-scale bases on the interval $[0, 1]$, using B-splines for the primal bases and the dual components from [5] adapted to the interval [10]. Tensor products yield corresponding dual pairs on \square . Using the parametric liftings γ_i and gluing across patch boundaries leads to globally continuous single-scale bases $\Phi_j, \tilde{\Phi}_j$ on Γ [3, 6, 14, 19]. For B-splines of order d and duals of order $\tilde{d} \geq d$ such that $d + \tilde{d}$ is even, the $\Phi_j, \tilde{\Phi}_j$ have approximation orders d, \tilde{d} , respectively. It is known that the respective regularity indices $\gamma, \tilde{\gamma}$ (inside each patch) satisfy $\gamma = d - 1/2$, while $\tilde{\gamma} > 0$ is known to increase proportionally to \tilde{d} . We refer the reader to [21] for a detailed description of the construction of wavelets on manifolds, including examples and figures.

In view of the biorthogonality of $\Phi_j, \tilde{\Phi}_j$, it will be convenient to employ the canonical projectors

$$(4.1) \quad Q_j v := \sum_{k \in \Delta_j} \langle v, \tilde{\phi}_{j,k} \rangle \phi_{j,k}, \quad Q_j^* v := \sum_{k \in \Delta_j} \langle v, \phi_{j,k} \rangle \tilde{\phi}_{j,k},$$

associated with the *multiresolution sequences* $\{V_j\}_{j>j_0}, \{\tilde{V}_j\}_{j>j_0}$. Here and below $j_0 + 1$ always stands for some fixed coarsest level of resolution that may depend on Γ .

It follows from the L^2 -boundedness of the Q_j that one has the following *Jackson*- and *Bernstein*-type estimates uniformly in j , namely,

$$(4.2) \quad \|v - Q_j v_j\|_s \lesssim 2^{-j(t-s)} \|v\|_t, \quad v \in H^t(\Gamma),$$

for all $-\tilde{d} \leq s \leq t \leq d, s < \gamma, -\tilde{\gamma} < t$ and

$$(4.3) \quad \|Q_j v\|_s \lesssim 2^{j(s-t)} \|Q_j v\|_t, \quad v \in H^t(\Gamma),$$

for all $t \leq s \leq \gamma$.

We introduce the index sets $\nabla_j := \Delta_{j+1} \setminus \Delta_j$. Given the single-scale bases $\Phi_j, \tilde{\Phi}_j$, one can construct now biorthogonal *complement bases* $\Psi_j = \{\psi_{j,k} : k \in \nabla_j\}, \tilde{\Psi}_j = \{\tilde{\psi}_{j,k} : k \in \nabla_j\}$, i.e., $\langle \psi_{j,k}, \tilde{\psi}_{j',k'} \rangle = \delta_{(j,k),(j',k')}$, such that

$$(4.4) \quad \text{diam supp } \psi_{j,k} \sim 2^{-j}, \quad j > j_0;$$

see, e.g., [3, 6, 13, 14] and [19] for particularly useful local representations of important construction ingredients. In fact, for these types of bases, the dual wavelets scale in the same way, but this will not be needed and does not hold for alternative constructions based on finite elements [15].

Denoting by W_j, \tilde{W}_j the span of Ψ_j , respectively, $\tilde{\Psi}_j$, biorthogonality implies that

$$V_{j+1} = W_j \oplus V_j, \quad \tilde{V}_{j+1} = \tilde{W}_j \oplus \tilde{V}_j, \quad \tilde{V}_j \perp W_j, \quad V_j \perp \tilde{W}_j.$$

Hence V_J and \tilde{V}_J can be written as a direct sum of the complement spaces W_j , respectively, $\tilde{W}_j, j_0 \leq j < J$ (using the convention $W_{j_0} := V_{j_0+1}, \tilde{W}_{j_0} := \tilde{V}_{j_0+1}, Q_{j_0} = Q_{j_0}^* := 0$). In fact, one has for $v_J \in V_J, \tilde{v}_J \in \tilde{V}_J$

$$v_J = \sum_{j=j_0}^{J-1} (Q_{j+1} - Q_j)v_J, \quad \tilde{v}_J = \sum_{j=j_0}^{J-1} (Q_{j+1}^* - Q_j^*)\tilde{v}_J,$$

where

$$(Q_{j+1} - Q_j)v = \sum_{k \in \nabla_j} \langle v, \tilde{\psi}_{j,k} \rangle \psi_{j,k}, \quad (Q_{j+1}^* - Q_j^*)v = \sum_{k \in \nabla_j} \langle v, \psi_{j,k} \rangle \tilde{\psi}_{j,k}.$$

A biorthogonal or *dual* pair of wavelet bases is now obtained by taking the coarse single-scale basis and the union of the complement bases $\Psi = \bigcup_{j \geq j_0} \Psi_j, \tilde{\Psi} = \bigcup_{j \geq j_0} \tilde{\Psi}_j$, where we have set for convenience $\Psi_{j_0} := \Phi_{j_0+1}, \tilde{\Psi}_{j_0} := \tilde{\Phi}_{j_0+1}$. We will refer to Ψ and $\tilde{\Psi}$ as the *primal*, respectively, *dual*, basis. Throughout the paper, all basis functions (scaling functions and wavelets) are normalized in $L^2(\Gamma)$.

From biorthogonality and the fact that the dual single-scale bases on \square represent all polynomials of order \tilde{d} exactly, one infers vanishing polynomial moments of the primal wavelets on \square , which, on account of the locality (4.4), entails the first key feature of the primal wavelets, namely, *vanishing moments* or the *cancellation property*

$$(4.5) \quad |\langle v, \psi_{j,k} \rangle| \lesssim 2^{-j(\tilde{d}+n/2)} |v|_{W^{\tilde{d},\infty}(\text{supp } \psi_{j,k})}.$$

Here $|v|_{W^{\tilde{d},\infty}(\Omega)} := \sup_{|\alpha|=\tilde{d}, x \in \Omega} |\partial^\alpha v(x)|$ denotes the seminorm in $W^{\tilde{d},\infty}(\Omega)$. The fact that the concept of biorthogonality allows us to choose the order \tilde{d} of vanishing moments higher than the approximation order d will be essential for deriving optimal compression strategies that could not be realized by orthonormal bases.

Of course, in the infinite-dimensional case the notion of basis has to be made more specific. The second key feature of the basis Ψ is the fact that (properly scaled versions of) $\Psi, \tilde{\Psi}$ are actually *Riesz bases* for a *whole range* of Sobolev spaces, i.e.,

$$(4.6) \quad \begin{aligned} \|v\|_t^2 &\sim \sum_{j \geq j_0} \sum_{k \in \nabla_j} 2^{2jt} |\langle v, \tilde{\psi}_{j,k} \rangle|^2, & t \in (-\tilde{\gamma}, \gamma), \\ \|v\|_t^2 &\sim \sum_{j \geq j_0} \sum_{k \in \nabla_j} 2^{2jt} |\langle v, \psi_{j,k} \rangle|^2, & t \in (-\gamma, \tilde{\gamma}). \end{aligned}$$

The validity of these norm equivalences hinges on the estimates (4.2) and (4.3) for *both* the primal and dual multiresolution sequences. The equivalences (4.6) will be essential for preconditioning.

5. Wavelet Galerkin schemes—preconditioning. As before let $A : H^q(\Gamma) \rightarrow H^{-q}(\Gamma)$ be a boundary integral operator of order $2q$. Since the wavelet basis Ψ is, in particular, a Riesz basis for $L^2(\Gamma)$, the associated system matrices

$$\mathbf{A}_J = [\langle A\psi_{j',k'}, \psi_{j,k} \rangle]_{j_0 \leq j, j' < J, k \in \nabla_j, k' \in \nabla_{j'}}$$

become more and more ill-conditioned when J increases. In fact, one has $\text{cond}_{l_2} \mathbf{A}_J \sim 2^{2J|q|}$. However, as a consequence of the stability of the Galerkin discretization under the given circumstances and the norm equivalences (4.6), the following simple *diagonal preconditioner* gives rise to uniformly bounded spectral condition numbers [8, 9, 11].

THEOREM 5.1. *Let the diagonal matrix \mathbf{D}_J^r be defined by*

$$[\mathbf{D}_J^r]_{(j,k),(j',k')} = 2^{rj} \delta_{(j,k),(j',k')}, \quad k \in \nabla_j, \quad k' \in \nabla_{j'}, \quad j_0 \leq j, j' < J.$$

If $A : H^q(\Gamma) \rightarrow H^{-q}(\Gamma)$ is a boundary integral operator of order $2q$, satisfying the assumptions (1), (2) from section 3, and if $\tilde{\gamma} > -q$, the diagonal matrix \mathbf{D}_J^{2q} defines an asymptotically optimal preconditioner for \mathbf{A}_J , i.e., $\text{cond}_{l_2}(\mathbf{D}_J^{-q} \mathbf{A}_J \mathbf{D}_J^{-q}) \sim 1$.

Although the above scaling is asymptotically optimal, it should be stressed that the quantitative performance may vary significantly among different scalings with the same asymptotic behavior. In particular, since Ψ is, on account of the mapping properties of A and the norm equivalences (4.6), also a Riesz basis with respect to the energy norm, it would be natural to normalize the wavelets in the energy norm which would suggest the specific scaling $\langle A\psi_{j,k}, \psi_{j,k} \rangle \sim 2^{2qj}$. In fact, this latter diagonal scaling improves and simplifies the wavelet preconditioning.

In view of the above simple preconditioning, the iterative solution of the Galerkin systems is feasible and its overall efficiency relies now on the cost of matrix/vector multiplications, which brings us to the central issue, namely, *matrix compression*.

6. Basic estimates. The basic ingredients in the analysis of the compression procedure are estimates for the matrix entries $\langle A\psi_{j',k'}, \psi_{j,k} \rangle$ with $k \in \nabla_j, k' \in \nabla_{j'}$ and $j, j' \geq j_0$. The convex hulls of the supports of the wavelets will be denoted by

$$(6.1) \quad \Omega_{j,k} := \text{conv hull}(\text{supp } \psi_{j,k}).$$

A complete proof of the following estimates can be found, e.g., in [15, 27].

THEOREM 6.1. *Suppose $n + 2\tilde{d} + 2q > 0$ and $j, j' > j_0$. Then one has*

$$|\langle A\psi_{j',k'}, \psi_{j,k} \rangle| \lesssim 2^{-(j+j')(\tilde{d}+n/2)} \text{dist}(\Omega_{j,k}, \Omega_{j',k'})^{-(n+2q+2\tilde{d})}$$

uniformly with respect to J .

However, in order to arrive ultimately at solution schemes with linear complexity, the number of nonzero entries in the compressed matrices should remain proportional to their size while preserving discretization error accuracy. To achieve this, it is not sufficient to consider only coefficients where the supports of the involved wavelets do not overlap. There are still $\mathcal{O}(N_J \log N_J) = \mathcal{O}(2^{Jn} J)$ coefficients that would remain. To avoid the logarithmic term we propose an additional, so-called *second compression*. For this purpose, we require that our primal basis functions be piecewise polynomial, in the sense that $\psi_{j,k}|_{\Gamma_{i,j+1,l}} = p \circ \gamma_i^{-1}$, where p is a polynomial. By

$$(6.2) \quad \Omega'_{j,k} := \text{sing supp } \psi_{j,k}$$

we denote the *singular support* of $\psi_{j,k}$, which is that subset of Γ where the function $\psi_{j,k}$ is not smooth. Thus the singular support of the wavelet $\psi_{j,k}$ consists of the boundaries of some of the elements $\Gamma_{i,j+1,l}$. The goal of the subsequent investigation is to estimate those matrix entries for which $\text{dist}(\Omega_{j,k}, \Omega'_{j',k'})$, $j \geq j'$, is sufficiently large.

To this end, we require the following extension lemma which follows, e.g., immediately from the well-known extension theorem of Calderón [28].

LEMMA 6.2. *The function $f_{i,j,k,l}$, defined by*

$$f_{i,j,k,l} := \psi_{j,k}|_{\Gamma_{i,j+1,l}} \circ \gamma_i = (\psi_{j,k} \circ \gamma_i)|_{C_{j+1,l}} \in C^\infty(C_{j+1,l}),$$

can be extended to a function $\tilde{f}_{i,j,k,l} \in C_0^\infty(\mathbb{R}^n)$ in such a way that $\text{diam supp } \tilde{f}_{i,j,k,l} \lesssim 2^{-j}$, $\tilde{f}_{i,j,k,l} \equiv \psi_{j,k} \circ \gamma_i$ on $C_{j+1,l}$, and that for all $s \geq 0$ there holds $\|\tilde{f}_{i,j,k,l}\|_{H^s(\mathbb{R}^n)} \lesssim 2^{js}$, independently of i, j, k, l .

Proof. Suppose that $f_\square \in C^\infty(\square)$ with $\|f_\square\|_{H^s(\square)} \lesssim 1$. By virtue of Calderón’s extension theorem, there exists an extension $f \in C_0^\infty(\mathbb{R}^n)$, i.e., $f(x) \equiv f_\square(x)$ on \square , satisfying $\|f\|_{H^s(\mathbb{R}^n)} \lesssim \|f_\square\|_{H^s(\square)}$. Let us consider an affine map κ with $\kappa(C_{j+1,l}) = \square$ and choose $f_{i,j,k,l}(x) := f_\square(\kappa(x))$. The claim follows now from $|\partial_{x_i} \kappa| = 2^{j+1}$, $i = 1, \dots, n$. \square

It is well known that boundary integral operators A of order $2q$, acting on smooth surfaces, are classical *pseudodifferential operators* [28]. Since the patches Γ_i are smooth and have smooth extensions $\tilde{\Gamma}_i$, there exists for each i a pseudodifferential operator $A^\sharp : H^q(\mathbb{R}^n) \rightarrow H^{-q}(\mathbb{R}^n)$ such that

$$(6.3) \quad A^\sharp f(x) = \int_{\mathbb{R}^n} \chi(x)\chi(y)k(\tilde{\gamma}_i(x), \tilde{\gamma}_i(y))f(y)\sqrt{\det \tilde{K}_i(y)}dy,$$

where χ is a C^∞ -cut-off function with respect to \square , i.e., $\chi(x) = 1$ on \square and $\chi(x) = 0$ outside $[-1, 2]^n$. Therefore $A^\sharp f(x) = A(f \circ \gamma_i)(\gamma_i(x))$ for all $f \in C_0^\infty(\square)$, $x \in \square$, and A^\sharp is compactly supported [29]. Moreover, it is well known [28] that the Schwartz kernel of pseudodifferential operators satisfies the standard estimate (2.6).

A compactly supported pseudodifferential operator $A^\sharp : H^s(\mathbb{R}^n) \rightarrow H^{s-2q}(\mathbb{R}^n)$ of order $2q$ acts continuously on Sobolev spaces [28, 29]. Therefore, for any function

$\tilde{f}_{i,j,k,l} \in C_0^\infty(\mathbb{R}^n)$, satisfying $\text{diam supp } \tilde{f}_{i,j,k,l} \sim 2^{-j}$ and $\|\tilde{f}_{i,j,k,l}\|_{H^s(\mathbb{R}^n)} \lesssim 2^{js}$ for all $s \geq 0$, one has $A^\sharp \tilde{f}_{i,j,k,l} \in C_0^\infty(\mathbb{R}^n)$ with

$$(6.4) \quad \|A^\sharp \tilde{f}_{i,j,k,l}\|_{H^{s-2q}(\mathbb{R}^n)} \lesssim 2^{js}.$$

With these preparations at hand, we are able to formulate the following result.

THEOREM 6.3. *Suppose that $n + 2\tilde{d} + 2q > 0$ and $j' > j \geq j_0$. Then, the coefficients $\langle A\psi_{j',k'}, \psi_{j,k} \rangle$ and $\langle A\psi_{j,k}, \psi_{j',k'} \rangle$ satisfy*

$$|\langle A\psi_{j',k'}, \psi_{j,k} \rangle|, |\langle A\psi_{j,k}, \psi_{j',k'} \rangle| \lesssim 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} \text{dist}(\Omega'_{j,k}, \Omega_{j',k'})^{-(2q+\tilde{d})},$$

uniformly with respect to j , provided that

$$(6.5) \quad \text{dist}(\Omega'_{j,k}, \Omega_{j',k'}) \gtrsim 2^{-j'}.$$

Proof. We shall consider three cases.

(i) The first observation concerns an estimate for disjoint supports that will be applied several times.

LEMMA 6.4. *Suppose that $\Omega_{j,k} \cap \Omega_{j',k'} = \emptyset$ and that f is any function supported on $\Omega_{j,k}$ satisfying $|f(x)| \lesssim 2^{jn/2}$, $x \in \Omega_{j,k}$. Then one has*

$$(6.6) \quad |\langle A\psi_{j',k'}, f \rangle| \lesssim 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} \text{dist}(\Omega_{j,k}, \Omega_{j',k'})^{-(2q+\tilde{d})}.$$

To prove (6.6) note that our assumption implies $\text{dist}(\Omega'_{j,k}, \Omega_{j',k'}) = \text{dist}(\Omega_{j,k}, \Omega_{j',k'})$. On account of the cancellation property (4.5) of the wavelet bases and the decay property (2.6) of the kernel, we obtain

$$\begin{aligned} |A\psi_{j',k'}(x)| &= |\langle k(x, \cdot), \psi_{j',k'} \rangle| \lesssim 2^{-j'(\tilde{d}+n/2)} |k(x, \cdot)|_{W^\infty, \tilde{d}(\Omega_{j',k'})} \\ &\lesssim 2^{-j'(\tilde{d}+n/2)} \text{dist}(x, \Omega_{j',k'})^{-(n+2q+\tilde{d})} \end{aligned}$$

for all $x \in \text{supp } \psi_{j,k}$. Therefore, we conclude that

$$\begin{aligned} |\langle A\psi_{j',k'}, f \rangle| &\lesssim \|f\|_{L^\infty(\Gamma)} \int_{\Omega_{j,k}} |A\psi_{j',k'}(x)| d\Gamma_x \\ &\lesssim 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} \int_{\Omega_{j,k}} \text{dist}(x, \Omega_{j',k'})^{-(n+2q+\tilde{d})} d\Gamma_x \\ &\leq 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} \text{dist}(\Omega_{j,k}, \Omega_{j',k'})^{-(2q+\tilde{d})}, \end{aligned}$$

which proves the lemma. Of course, the same reasoning applies to the adjoint boundary integral operator A^* .

(ii) Next, we treat the case $\Omega_{j,k} \cap \Omega_{j',k'} \neq \emptyset$ and $\Omega_{j,k} \subset \Gamma_i$. By (6.5) we have $\Omega_{j',k'} \subset \Omega_{j,k}$ i.e., both wavelets are supported on the same patch. We infer from (6.5) that there exists an element $\Omega_{j',k'} \subset \Gamma_{i,j+1,l} \subset \Omega_{j,k}$ such that

$$f_{i,j,k,l} := \psi_{j,k}|_{\Gamma_{i,j+1,l}} \circ \gamma_i = (\psi_{j,k} \circ \gamma_i)|_{C_{j+1,l}}$$

is a $C^\infty(C_{j+1,l})$ function. On account of Lemma 6.2, we can choose an extension of $f_{i,j,k,l}$, denoted by $\tilde{f}_{i,j,k,l}$. Decomposing $\psi_{j,k} \circ \gamma_i = \tilde{f}_{i,j,k,l} + \tilde{f}_{i,j,k,l}^C$, we obtain

$$\begin{aligned} |\langle A\psi_{j,k}, \psi_{j',k'} \rangle| &= \left| \int_{\mathbb{R}^n} A^\sharp(\tilde{f}_{i,j,k,l} + \tilde{f}_{i,j,k,l}^C)(x)(\psi_{j',k'} \circ \gamma_i)(x) dx \right| \\ &\leq \left| \int_{\mathbb{R}^n} A^\sharp \tilde{f}_{i,j,k,l}(x)(\psi_{j',k'} \circ \gamma_i)(x) dx \right| \\ &\quad + \left| \int_{\mathbb{R}^n} A^\sharp \tilde{f}_{i,j,k,l}^C(x)(\psi_{j',k'} \circ \gamma_i)(x) dx \right|. \end{aligned}$$

The second term on the right-hand side can be treated analogously to (6.6), i.e.,

$$\left| \int_{\mathbb{R}^n} A^\sharp \tilde{f}_{i,j,k,l}^C(x)(\psi_{j',k'} \circ \gamma_i)(x) dx \right| \lesssim 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} \text{dist}(\Omega'_{j,k}, \Omega_{j',k'})^{-(2q+\tilde{d})},$$

because $\text{dist}(\text{supp} \tilde{f}_{i,j,k,l}^C, \text{supp}(\psi_{j',k'} \circ \gamma_i)) \sim \text{dist}(\Omega'_{j,k}, \Omega_{j',k'})$. Invoking (4.5) and (6.4), the first term can be estimated by

$$\left| \int_{\mathbb{R}^n} A^\sharp \tilde{f}_{i,j,k,l}(x)(\psi_{j',k'} \circ \gamma_i)(x) dx \right| \lesssim 2^{-j'(\tilde{d}+n/2)} \|A^\sharp \tilde{f}_{i,j,k,l}(x)\|_{W^{\infty, \tilde{d}}(\text{supp}(\psi_{j',k'} \circ \gamma_i))}.$$

By virtue of Sobolev’s embedding theorem, this implies, in view of (6.4),

$$\begin{aligned} \left| \int_{\mathbb{R}^n} A^\sharp \tilde{f}_{i,j,k,l}(x)(\psi_{j',k'} \circ \gamma_i)(x) dx \right| &\lesssim 2^{-j'(\tilde{d}+n/2)} \|A^\sharp \tilde{f}_{i,j,k,l}(x)\|_{H^{\tilde{d}+n/2}(\mathbb{R}^n)} \\ &\lesssim 2^{-j'(\tilde{d}+n/2)} 2^{j(\tilde{d}+2q+n/2)}. \end{aligned}$$

Since $\Omega_{j,k} \cap \Omega_{j',k'} \neq \emptyset$, one has, in view of (6.5) and $j' \geq j$, that $\text{dist}(\Omega'_{j,k}, \Omega_{j',k'}) \lesssim 2^{-j}$, so that we arrive at the desired estimate

$$\left| \int_{\mathbb{R}^n} A^\sharp \tilde{f}_{i,j,k,l}(x)(\psi_{j',k'} \circ \gamma_i)(x) dx \right| \lesssim 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} \text{dist}(\Omega'_{j,k}, \Omega_{j',k'})^{-(2q+\tilde{d})}.$$

(iii) It remains to consider the case $\Omega_{j,k} \cap \Omega_{j',k'} \neq \emptyset$, where, however, $\psi_{j,k}$ is not supported completely in the patch Γ_i . In this case, we decompose $\psi_{j,k} = (\psi_{j,k} - \psi_{j,k}|_{\Gamma_i}) + \psi_{j,k}|_{\Gamma_i}$. Invoking (6.6), we derive

$$|\langle A(\psi_{j,k} - \psi_{j,k}|_{\Gamma_i}), \psi_{j',k'} \rangle| \lesssim 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} \text{dist}(\Omega'_{j,k}, \Omega_{j',k'})^{-(2q+\tilde{d})}$$

because we have again $\text{dist}(\text{supp}(\psi_{j,k} - \psi_{j,k}|_{\Gamma_i}), \Omega_{j',k'}) \geq \text{dist}(\Omega'_{j,k}, \Omega_{j',k'})$. Finally, estimating $|\langle A(\psi_{j,k}|_{\Gamma_i}), \psi_{j',k'} \rangle|$ as in step (ii) finishes the proof. \square

Remark. We recall from [8, 27] that there is a general estimate which states that the matrix entries for wavelets with overlapping supports decay with increasing difference of scales. In fact, for each $0 \leq \delta < \gamma - q$ we have $|\langle A\psi_{j',k'}, \psi_{j,k} \rangle| \lesssim 2^{-\delta|j-j'|}$. Since $\gamma < d$ this estimate is, however, not sufficient to achieve the optimal order of convergence within the desired linear complexity.

7. Matrix compression. The discretization of a boundary integral operator $A : H^q(\Gamma) \rightarrow H^{-q}(\Gamma)$ by wavelets with a sufficiently strong cancellation property (4.5) yields, in view of the above estimates, quasi-sparse matrices. In the first compression step all matrix entries, for which the distance of the supports of the corresponding trial and test functions is larger than a level depending cut-off parameter $\mathcal{B}_{j,j'}$, are set to zero. In the second compression step also some of those matrix entries are neglected, for which the corresponding trial and test functions have overlapping supports.

A priori compression. Let $\Omega_{j,k}$ and $\Omega'_{j,k}$ be given as in (6.1) and (6.2). Then, the compressed system matrix \mathbf{A}_J^ϵ , corresponding to the boundary integral operator A , is defined by

$$(7.1) \quad [\mathbf{A}_J^\epsilon]_{(j,k),(j',k')} := \begin{cases} 0, & \text{dist}(\Omega_{j,k}, \Omega'_{j',k'}) > \mathcal{B}_{j,j'} \text{ and } j, j' > j_0, \\ 0, & \text{dist}(\Omega_{j,k}, \Omega'_{j',k'}) \lesssim 2^{-\min\{j,j'\}} \text{ and} \\ & \text{dist}(\Omega'_{j,k}, \Omega_{j',k'}) > \mathcal{B}'_{j,j'} \text{ if } j' > j \geq j_0, \\ & \text{dist}(\Omega_{j,k}, \Omega'_{j',k'}) > \mathcal{B}'_{j,j'} \text{ if } j > j' \geq j_0, \\ \langle A\psi_{j',k'}, \psi_{j,k} \rangle, & \text{otherwise.} \end{cases}$$

Fixing

$$(7.2) \quad a, a' > 1, \quad d < d' < \tilde{d} + 2q,$$

the cut-off parameters $\mathcal{B}_{j,j'}$ and $\mathcal{B}'_{j,j'}$ are set as follows:

$$(7.3) \quad \begin{aligned} \mathcal{B}_{j,j'} &= a \max \left\{ 2^{-\min\{j,j'\}}, 2^{\frac{2J(d'-q)-(j+j')(d'+\tilde{d})}{2(d+q)}} \right\}, \\ \mathcal{B}'_{j,j'} &= a' \max \left\{ 2^{-\max\{j,j'\}}, 2^{\frac{2J(d'-q)-(j+j')d'-\max\{j,j'\}\tilde{d}}{d+2q}} \right\}. \end{aligned}$$

Remark 7.1. Relation (7.2) requires the order of vanishing moments, viz. the order of exactness of the dual basis, to exceed the order of the primal basis by an amount determined by the order of the operator $d < \tilde{d} + 2q$. In the case of equality the matrices are still compressible, but additional log terms arise in the complexity estimates. One can find many pairs of biorthogonal wavelets from the family in [14, 21] satisfying this relation. To obtain quantitatively best performance we employ those with possibly small support, i.e., with possibly small d satisfying $d < \tilde{d} + 2q$. This choice is confirmed by numerical experience.

The parameter a is a fixed constant which determines the bandwidth in the block matrices $\mathbf{A}_{j,j'}^\epsilon := [\mathbf{A}_J^\epsilon]_{(j,\nabla_j),(j',\nabla_{j'})}$, $j_0 \leq j, j' < J$. We emphasize that the parameters a and a' are independent of J .

When the entries of the compressed system matrix \mathbf{A}_J^ϵ have been computed, we apply an *a posteriori compression* by setting all entries to zero, which are smaller than a level-dependent threshold. In this way, a matrix $\tilde{\mathbf{A}}_J^\epsilon$ is obtained which has even less nonzero entries than the matrix \mathbf{A}_J^ϵ . Although this does not accelerate the computation of the matrix coefficients, the amount of necessary memory for storing the system matrix is reduced considerably.

A posteriori compression. We define the a posteriori compression by

$$(7.4) \quad [\tilde{\mathbf{A}}_J^\epsilon]_{(j,k),(j',k')} = \begin{cases} 0 & \text{if } |[\mathbf{A}_J^\epsilon]_{(j,k),(j',k')}| \leq \varepsilon_{j,j'}, \\ [\mathbf{A}_J^\epsilon]_{(j,k),(j',k')} & \text{if } |[\mathbf{A}_J^\epsilon]_{(j,k),(j',k')}| > \varepsilon_{j,j'}. \end{cases}$$

Here the level-dependent threshold $\varepsilon_{j,j'}$ is chosen as

$$(7.5) \quad \varepsilon_{j,j'} = a'' \min \left\{ 2^{-\lfloor \frac{j-j'}{2} \rfloor n}, 2^{-n(J-\frac{j+j'}{2})\frac{d'-q}{d+q}} \right\} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}$$

with $a'' < 1$ and $d' \in (d, \tilde{d} + 2q)$ from (7.2).

8. Matrix estimates. In order to study the accuracy of the solutions to the compressed systems, we investigate the perturbation introduced by discarding specific matrix elements. The perturbation matrices are of scalewise blocks of the type $\mathbf{R}_{j,j'} := \mathbf{A}_{j,j'} - \mathbf{A}_{j,j'}^\epsilon$. By $\|\mathbf{R}_{j,j'}\|_p$ we denote the operator norm of the matrix $\mathbf{R}_{j,j'}$ with respect to the norm l^p .

In order to analyze the error introduced by our compression strategy, we decompose the complete compression into three subsequent steps.

THEOREM 8.1 (first compression). *We define the matrix $\mathbf{A}_j^{\epsilon_1}$ by*

$$[\mathbf{A}_j^{\epsilon_1}]_{(j,k),(j',k')} := \begin{cases} 0, & \text{dist}(\Omega_{j,k}, \Omega_{j',k'}) > \mathcal{B}_{j,j'} \text{ and } j, j' > j_0, \\ \langle A\psi_{j',k'}, \psi_{j,k} \rangle, & \text{otherwise.} \end{cases}$$

Here the parameter $\mathcal{B}_{j,j'}$ is given by (7.2) and (7.3). Then, one has for the perturbation matrix $\mathbf{R}_{j,j'} := \mathbf{A}_{j,j'} - \mathbf{A}_{j,j'}^{\epsilon_1}$

$$\|\mathbf{R}_{j,j'}\|_2 \lesssim a^{-2(\tilde{d}+q)} 2^{2Jq} 2^{-2d'(J - \frac{j+j'}{2})}.$$

Proof. We proceed in two steps.

(i) We abbreviate $\mathbf{R}_{j,j'} := [r_{(j,k),(j',k')}]_{k \in \nabla_j, k' \in \nabla_{j'}}$. Invoking Theorem 6.1, we find for the column sum

$$\begin{aligned} \sum_{k \in \nabla_j} |r_{(j,k),(j',k')}| &= \sum_{\{k \in \nabla_j : \text{dist}(\Omega_{j,k}, \Omega_{j',k'}) > \mathcal{B}_{j,j'}\}} |\langle A\psi_{j',k'}, \psi_{j,k} \rangle| \\ &\lesssim \sum_{\{k \in \nabla_j : \text{dist}(\Omega_{j,k}, \Omega_{j',k'}) > \mathcal{B}_{j,j'}\}} \\ &\quad \times 2^{-(j+j')(\tilde{d}+n/2)} \text{dist}(\Omega_{j,k}, \Omega_{j',k'})^{-(n+2\tilde{d}+2q)}. \end{aligned}$$

Since $\mathcal{B}_{j,j'} \geq a \max\{2^{-j}, 2^{-j'}\}$, we can estimate this sum by an integral which yields

$$\begin{aligned} \sum_{k \in \nabla_j} |r_{(j,k),(j',k')}| &\lesssim 2^{-(j+j')(\tilde{d}+n/2)} 2^{jn} \int_{\|x\| > \mathcal{B}_{j,j'}} \|x\|^{-(n+2\tilde{d}+2q)} dx \\ &\lesssim 2^{-(j+j')(\tilde{d}+n/2)} 2^{jn} \mathcal{B}_{j,j'}^{-2(\tilde{d}+q)}. \end{aligned}$$

On the other hand, inserting the estimate $\mathcal{B}_{j,j'} \geq a 2^{\frac{2J(d'-q)-(j+j')(\tilde{d}+d')}{2(d+q)}}$ (see (7.3)), we arrive at

$$\sum_{k \in \nabla_j} |r_{(j,k),(j',k')}| \lesssim a^{-2(\tilde{d}+q)} 2^{\frac{(j-j')n}{2}} 2^{2Jq} 2^{-2d'(J - \frac{j+j'}{2})}.$$

In complete analogy, one proves an analogous estimate for the row sums,

$$\sum_{k' \in \nabla_{j'}} |r_{(j,k),(j',k')}| \lesssim a^{-2(\tilde{d}+q)} 2^{\frac{(j'-j)n}{2}} 2^{2Jq} 2^{-2d'(J - \frac{j+j'}{2})}.$$

(ii) From the estimate for the operator norms of matrices $\|\mathbf{R}_{j,j'}\|_2^2 \leq \|\mathbf{R}_{j,j'}\|_1 \|\mathbf{R}_{j,j'}\|_\infty$, it is easy to conclude the following version of the Schur lemma (see, e.g., [23, 27]):

$$\begin{aligned} \|\mathbf{R}_{j,j'}\|_2 &\leq \left[\max_{k \in \nabla_j} \sum_{k' \in \nabla_{j'}} 2^{\frac{(j-j')n}{2}} |r_{(j,k),(j',k')}| \right]^{1/2} \left[\max_{k' \in \nabla_{j'}} \sum_{k \in \nabla_j} 2^{\frac{(j'-j)n}{2}} |r_{(j,k),(j',k')}| \right]^{1/2} \\ &\lesssim a^{-2(\tilde{d}+q)} 2^{2Jq} 2^{-2d'(J - \frac{j+j'}{2})}, \end{aligned}$$

which proves the assertion. \square

The following so-called second compression concerns entries involving basis functions with overlapping supports. It is important that here the coarse scale basis function may be a scaling function which greatly affects the near field compression.

THEOREM 8.2 (second compression). *In addition to the first compression we apply the following second compression:*

$$[\mathbf{A}_J^{\epsilon_2}]_{(j,k),(j',k')} := \begin{cases} 0, & \text{dist}(\Omega_{j,k}, \Omega_{j',k'}) \lesssim 2^{-\min\{j,j'\}} \text{ and} \\ & \text{dist}(\Omega'_{j,k}, \Omega'_{j',k'}) > \mathcal{B}'_{j,j'} \text{ if } j' > j \geq j_0, \\ & \text{dist}(\Omega_{j,k}, \Omega'_{j',k'}) > \mathcal{B}'_{j,j'} \text{ if } j > j' \geq j_0, \\ [\mathbf{A}_J^{\epsilon_1}]_{(j,k),(j',k')}, & \text{otherwise,} \end{cases}$$

where the parameter $\mathcal{B}'_{j,j'}$ is set in accordance with (7.2) and (7.3). Then, the corresponding perturbation matrix $\mathbf{S}_{j,j'} := \mathbf{A}_{j,j'}^{\epsilon_1} - \mathbf{A}_{j,j'}^{\epsilon_2}$ satisfies

$$\|\mathbf{S}_{j,j'}\|_2 \lesssim (a')^{-(\tilde{d}+2q)} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}.$$

Proof. Abbreviating $\mathbf{S}_{j,j'} := [s_{(j,k),(j',k')}]_{k \in \nabla_j, k' \in \nabla_{j'}}$ and assuming without loss of generality that $j' > j$, we infer from Theorem 6.3 that

$$\begin{aligned} |s_{(j,k),(j',k')}| &\lesssim 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} \mathcal{B}_{j,j'}^{-(2q+\tilde{d})} \\ &\lesssim (a')^{-(\tilde{d}+2q)} 2^{jn/2} 2^{-j'(\tilde{d}+n/2)} 2^{-2J(d'-q)+(j+j')d'+j'\tilde{d}} \\ &= (a')^{-(\tilde{d}+2q)} 2^{(j-j')n/2} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}. \end{aligned}$$

The condition $\text{dist}(\Omega_{j,k}, \Omega_{j',k'}) \lesssim 2^{-\min\{j,j'\}}$ guarantees that in each row and column of $\mathbf{S}_{j,j'}$ we have set at most $\mathcal{O}(2^{(j'-j)n})$, respectively, $\mathcal{O}(1)$ entries to zero. Therefore, we obtain for the weighted row sums

$$\begin{aligned} \sum_{k \in \nabla_j} 2^{\frac{(j-j')n}{2}} |s_{(j,k),(j',k')}| &\lesssim \sum_{k' \in \nabla_{j'}} (a')^{-(\tilde{d}+2q)} 2^{j'n} 2^{(j-j')n} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})} \\ &\lesssim (a')^{-(\tilde{d}+2q)} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}, \end{aligned}$$

and likewise for the weighted column sums

$$\sum_{k' \in \nabla_{j'}} 2^{\frac{(j'-j)n}{2}} |s_{(j,k),(j',k')}| \lesssim (a')^{-(\tilde{d}+2q)} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}$$

for all $j_0 \leq j < j' < J$. In complete analogy to the proof of Theorem 8.1 we conclude the assertion. \square

THEOREM 8.3 (a posteriori compression). *Let the matrix \mathbf{A}_J^ϵ be compressed according to Theorems 8.1 and 8.2. Then the a posteriori compression defined by (7.4) with the level-dependent threshold $\varepsilon_{j,j'}$ from (7.5) causes a block perturbation $\mathbf{T}_{j,j'} := \tilde{\mathbf{A}}_{j,j'}^\epsilon - \mathbf{A}_{j,j'}^\epsilon$ satisfying*

$$\|\mathbf{T}_{j,j'}\|_2 \lesssim a'' 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}.$$

Proof. We organize the proof in four steps.

(i) Abbreviating $\mathbf{T}_{j,j'} := [t_{(j,k),(j',k')}]_{k \in \nabla_j, k' \in \nabla_{j'}}$, one obviously has

$$(8.1) \quad |t_{(j,k),(j',k')}| \leq a'' \min \left\{ 2^{-\frac{|j-j'|n}{2}}, 2^{-n(J-\frac{j+j'}{2})\frac{d'-q}{d+q}} \right\} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}.$$

We shall use the first compression in order to derive from this inequality the desired. To this end, we find in each row and column of $\mathbf{T}_{j,j'}$ only $\mathcal{O}([\mathcal{B}_{j,j'} 2^{j'}]^n)$, respectively, $\mathcal{O}([\mathcal{B}_{j,j'} 2^j]^n)$ nonzero entries. Setting $M := \frac{d'+\tilde{d}}{2(d+q)}$, one has

$$2^{\frac{2J(d'-q)-(j+j')(d'+\tilde{d})}{2(d+q)}} = 2^{-J} 2^{\frac{(j-j)(d'+\tilde{d})}{2(d+q)}} 2^{\frac{(j-j')(d'+\tilde{d})}{2(d+q)}} = 2^{-J} 2^{(J-j)M} 2^{(J-j')M}.$$

Hence, by (7.3), the cut-off parameter for the first compression takes the form

$$(8.2) \quad \mathcal{B}_{j,j'} \sim \max \left\{ 2^{-\min\{j,j'\}}, 2^{-J} 2^{(J-j)M} 2^{(J-j')M} \right\}.$$

From (7.2) and $q < d - \frac{1}{2}$, one concludes $\frac{1}{2} < M < 1$. Moreover, we shall make use of the identity

$$(8.3) \quad 2^{-n(J-\frac{j+j'}{2})\frac{d'-q}{d+q}} = 2^{-2n(J-\frac{j+j'}{2})(M-\frac{1}{2})}.$$

Without loss of generality, we assume in the following that $j' \geq j$.

(ii) With these preparations at hand we shall first estimate the block matrices $\mathbf{T}_{j,j'}$ with $2^{-\frac{(j'-j)n}{2}} \leq 2^{-2n(J-\frac{j+j'}{2})(M-\frac{1}{2})}$. One readily verifies that this relation is equivalent to $2^{-j} \geq 2^{-J} 2^{(J-j)M} 2^{(J-j')M}$, which, by (8.2), implies that the cut-off parameter satisfies $\mathcal{B}_{j,j'} \sim 2^{-j}$. Thus, from (8.1) one infers the estimate

$$\begin{aligned} \sum_{k' \in \nabla_{j'}} 2^{\frac{(j-j')n}{2}} |t_{(j,k),(j',k')}| &\lesssim a'' 2^{\frac{(j+j')n}{2}} 2^{-jn} 2^{-\frac{(j'-j)n}{2}} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})} \\ &= a'' 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})} \end{aligned}$$

for the weighted row sums of $\mathbf{T}_{j,j'}$. Analogously, one derives

$$\sum_{k \in \nabla_j} 2^{\frac{(j'-j)n}{2}} |t_{(j,k),(j',k')}| \lesssim a'' 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}$$

for the weighted column sums.

(iii) We still have to estimate the errors in the remaining blocks, where $2^{-\frac{(j'-j)n}{2}} > 2^{-2n(J-\frac{j+j'}{2})(M-\frac{1}{2})}$. Then, by (8.3), the cut-off parameter is given by $\mathcal{B}_{j,j'} \sim 2^{-J} 2^{(J-j)M} 2^{(J-j')M}$. Therefore, we obtain for the weighted row sums

$$\begin{aligned} \sum_{k' \in \nabla_{j'}} 2^{\frac{(j-j')n}{2}} |t_{(j,k),(j',k')}| &\lesssim a'' 2^{\frac{(j+j')n}{2}} 2^{-Jn} 2^{(J-j)Mn} 2^{(J-j')Mn} 2^{-2n(J-\frac{j+j'}{2})(M-\frac{1}{2})} 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})} \\ &= a'' 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}, \end{aligned}$$

and a similar estimate for the weighted column sums.

(iv) Combining the estimates in steps (ii) and (iii), we conclude that

$$\sum_{k' \in \nabla_{j'}} 2^{\frac{(j-j')n}{2}} |t_{(j,k),(j',k')}|, \sum_{k \in \nabla_j} 2^{\frac{(j'-j)n}{2}} |t_{(j,k),(j',k')}| \lesssim a'' 2^{2Jq} 2^{-2d'(J-\frac{j+j'}{2})}$$

for all $j_0 \leq j, j' < J$. The proof can now be completed in complete analogy to the proof of Theorem 8.1. \square

9. Consistency. We shall establish next the consistency of the compressed scheme with the original operator equation in the corresponding Sobolev norms. To this end, note that the operator $\tilde{A}_J^\epsilon : H^s(\Gamma) \rightarrow H^{s-2q}(\Gamma)$, $-\tilde{\gamma} < s < \tilde{\gamma} + 2q$, defined by

$$A_J^\epsilon u = \sum_{j,j'=j_0}^{J-1} \sum_{k,k'} \tilde{\psi}_{j,k} [\tilde{\mathbf{A}}_J^\epsilon]_{(j,k),(j',k')} \langle \tilde{\psi}_{j,k}, u \rangle,$$

is represented by the compressed system matrix, since apparently $\langle \tilde{A}_J^\epsilon \psi_{j',k'}, \psi_{j,k} \rangle = [\tilde{\mathbf{A}}_J^\epsilon]_{(j,k),(j',k')}$.

THEOREM 9.1. *Let $d < \tilde{d} + 2q$ and $\tilde{\mathbf{A}}_J^\epsilon$ be the compressed matrix, defined according to section 7. Then, for $q \leq t, t' \leq d$ the estimate*

$$(9.1) \quad | \langle (A - \tilde{A}_J^\epsilon) Q_J u, Q_J v \rangle | \lesssim \epsilon 2^{J(2q-t-t')} \|u\|_t \|v\|_{t'}$$

holds uniformly with respect to J , where

$$(9.2) \quad \epsilon := a^{-2(\tilde{d}+q)} + (a')^{-(\tilde{d}+2q)} + a'',$$

and a, a', a'' are the constants from (7.2) and (7.5).

Proof. By definition of the block perturbation matrices $\mathbf{R}_{j,j'}$, $\mathbf{S}_{j,j'}$, $\mathbf{T}_{j,j'}$, one has

$$| \langle (A - \tilde{A}_J^\epsilon) \psi_{j',k'}, \psi_{j,k} \rangle | \leq | [\mathbf{R}_{j,j'} + \mathbf{S}_{j,j'} + \mathbf{T}_{j,j'}]_{k,k'} |.$$

Hence, we can estimate

$$\begin{aligned} (9.3) \quad & | \langle (A - \tilde{A}_J^\epsilon) Q_J u, Q_J v \rangle | \\ &= \left| \sum_{j,j'=j_0}^{J-1} \langle (A - \tilde{A}_J^\epsilon) (Q_{j'+1} - Q_{j'}) u, (Q_{j+1} - Q_j) v \rangle \right| \\ &\leq \sum_{j,j'=j_0}^{J-1} | \langle (A - \tilde{A}_J^\epsilon) (Q_{j'+1} - Q_{j'}) u, (Q_{j+1} - Q_j) v \rangle | \\ &= \sum_{j,j'=j_0}^{J-1} \left| \sum_{k \in \nabla_j} \sum_{k' \in \nabla_{j'}} \langle (A - \tilde{A}_J^\epsilon) \psi_{j',k'}, \psi_{j,k} \rangle \langle u, \tilde{\psi}_{j',k'} \rangle \langle v, \tilde{\psi}_{j,k} \rangle \right| \\ &\leq \sum_{j,j'=j_0}^{J-1} \| \mathbf{R}_{j,j'} + \mathbf{S}_{j,j'} + \mathbf{T}_{j,j'} \|_2 \| [\langle u, \tilde{\psi}_{j',k'} \rangle]_{k' \in \nabla_{j'}} \|_2 \| [\langle v, \tilde{\psi}_{j,k} \rangle]_{k \in \nabla_j} \|_2. \end{aligned}$$

Invoking the inverse estimate (4.3) and the approximation property (4.2) yields

$$\begin{aligned} \| [\langle u, \tilde{\psi}_{j',k'} \rangle]_{k' \in \nabla_{j'}} \|_2 &\sim \| (Q_{j'+1} - Q_{j'}) u \|_0 \lesssim 2^{-j't} \|u\|_t, \\ \| [\langle v, \tilde{\psi}_{j,k} \rangle]_{k \in \nabla_j} \|_2 &\sim \| (Q_{j+1} - Q_j) v \|_0 \lesssim 2^{-jt'} \|v\|_{t'}. \end{aligned}$$

Further, from Theorems 8.1, 8.2, and 8.3, we conclude

$$\| \mathbf{R}_{j,j'} + \mathbf{S}_{j,j'} + \mathbf{T}_{j,j'} \|_2 \lesssim \epsilon 2^{2Jq-2d'(J-\frac{j+j'}{2})}.$$

Inserting these estimates in (9.3) provides

$$\begin{aligned} |\langle (A - \tilde{A}_J^\epsilon)Q_J u, Q_J v \rangle| &\lesssim \epsilon 2^{J(2q-t-t')} \|u\|_t \|v\|_{t'} \sum_{j, j'=j_0}^{J-1} 2^{-j'(d'-t)} 2^{-j(d'-t')} \\ &\lesssim \epsilon 2^{J(2q-t-t')} \|u\|_t \|v\|_{t'} \end{aligned}$$

since $t, t' \leq d < d'$. \square

10. Convergence. With the estimates of section 8 at hand we can prove that the proposed compression strategy retains the optimal order of convergence of the underlying Galerkin scheme. In this context, we shall encounter conditions on the parameters a, a', a'' defining ϵ in (9.2). From Theorem 9.1 we deduce

$$|\langle (A - \tilde{A}_J^\epsilon)u_J, u_J \rangle| \leq \epsilon \|u_J\|_q^2,$$

which implies the V_J ellipticity. Indeed, inserting this result into (3.3), we get for $J > J_0$ that

$$|\langle (\tilde{A}_J^\epsilon + \tilde{A}_J^{\epsilon*})u_J, u_J \rangle| \geq (c - 2\epsilon) \|u_J\|_q^2 \gtrsim \|u_J\|_q^2,$$

with $c > 0$, if ϵ from (9.2) is sufficiently small.

THEOREM 10.1 (stability). *Let ϵ from (9.2) be sufficiently small. Then, the matrix $\tilde{\mathbf{A}}_J^\epsilon$, which arises by the compression according to (7.1) and (7.4), defines a stable scheme, i.e., $\|\tilde{A}_J^\epsilon u_J\|_{-q} \sim \|u_J\|_q$, uniformly in $J > J_0$.*

This theorem is an immediate consequence of Lemma 3.1 and the norm equivalences, which already requires that $\tilde{\gamma} > -q$. In the limit case $\tilde{\gamma} = -q$ a more sophisticated proof presented in [26] shows that Theorem 10.1 remains valid.

THEOREM 10.2 (convergence). *Let ϵ from (9.2) be sufficiently small to ensure uniform stability of \tilde{A}_J^ϵ . Then, the solution $u_J = \sum_{j=j_0}^{J-1} \sum_{k \in \nabla_j} u_{j,k} \psi_{j,k}$ of the compressed scheme $\tilde{\mathbf{A}}_J^\epsilon \mathbf{u}_J = \mathbf{f}_J$, where $\mathbf{u}_J = [u_{j,k}]_{j_0 \leq j < J, k \in \nabla_j}$, differs from the exact solution u , satisfying $Au = f$, in the energy norm only by*

$$\|u - u_J\|_q \lesssim 2^{J(q-d)} \|u\|_d$$

uniformly in J .

Proof. Strang’s first lemma [4] provides

$$\|u - u_J\|_q \lesssim \inf_{v_J \in V_J} \left\{ \|u - v_J\|_q + \sup_{w_J \in V_J} \frac{|\langle (A - \tilde{A}_J^\epsilon)v_J, w_J \rangle|}{\|w_J\|_q} \right\}.$$

The consistency (Theorem 9.1) implies that

$$|\langle (A - \tilde{A}_J^\epsilon)Q_J u, w_J \rangle| = |\langle (A - \tilde{A}_J^\epsilon)Q_J u, Q_J w_J \rangle| \lesssim 2^{J(q-d)} \|u\|_d \|w_J\|_q$$

for all $u \in H^d(\Gamma)$ and $w_J \in V_J$. Hence, choosing $v_J := Q_J u$, we arrive at

$$\begin{aligned} \|u - u_J\|_q &\lesssim \|u - Q_J u\|_q + \sup_{w_J \in V_J} \frac{|\langle (A - \tilde{A}_J^\epsilon)Q_J u, Q_J w_J \rangle|}{\|w_J\|_q} \\ &\lesssim 2^{J(q-d)} \|u\|_d. \quad \square \end{aligned}$$

THEOREM 10.3 (Aubin–Nitsche). *In addition to the assumptions of Theorem 10.2 suppose that $\|A^*v\|_{t-q} \sim \|v\|_{t+q}$ for all $0 \leq t \leq d - q$, i.e., $A^* : H^{t+q}(\Gamma) \rightarrow H^{t-q}(\Gamma)$ is an isomorphism. Then the error estimate*

$$\|u - u_J\|_{q-t} \lesssim 2^{J(q-d-t)} \|u\|_d$$

holds for all $0 \leq t \leq d - q$.

Proof. Recalling that

$$\|u - u_J\|_{q-t} = \sup_{g \in H^{t-q}(\Gamma)} \frac{\langle u - u_J, g \rangle}{\|g\|_{t-q}}.$$

we obtain for $v \in H^{t+q}(\Gamma)$ with $A^*v = g$

$$\|u - u_J\|_{q-t} = \sup_{v \in H^{t+q}(\Gamma)} \frac{|\langle A(u - u_J), v \rangle|}{\|v\|_{t+q}}.$$

Utilizing the Galerkin orthogonality $\langle \tilde{A}_J^\epsilon u_J, Q_J v \rangle = \langle Au, Q_J v \rangle$, we can decompose

$$\begin{aligned} \langle A(u - u_J), v \rangle &= \langle A(u - u_J), v - Q_J v \rangle + \langle A(u - u_J), Q_J v \rangle \\ &= \langle A(u - u_J), v - Q_J v \rangle - \langle (A - \tilde{A}_J^\epsilon) u_J, Q_J v \rangle. \end{aligned}$$

The first term on the right-hand side is estimated by Theorem 10.2 in combination with the approximation property (4.2),

$$|\langle A(u - u_J), v - Q_J v \rangle| \lesssim \|u - u_J\|_q \|v - Q_J v\|_q \lesssim 2^{J(q-d-t)} \|u\|_d \|v\|_{t+q}.$$

For the second term we obtain, on account of Theorem 9.1,

$$\begin{aligned} |\langle (A - \tilde{A}_J^\epsilon) u_J, Q_J v \rangle| &\leq |\langle (A - \tilde{A}_J^\epsilon)(u_J - Q_J u), Q_J v \rangle| + |\langle (A - \tilde{A}_J^\epsilon) Q_J u, Q_J v \rangle| \\ &\lesssim 2^{-Jt} \|u_J - Q_J u\|_q \|v\|_{t+q} + 2^{J(q-d-t)} \|u\|_d \|v\|_{t+q}. \end{aligned}$$

Inserting $\|u_J - Q_J u\|_q \leq \|u - u_J\|_q + \|u - Q_J u\|_q \lesssim 2^{J(q-d)} \|u\|_d$ yields

$$|\langle (A - \tilde{A}_J^\epsilon) u_J, Q_J v \rangle| \lesssim 2^{J(q-d-t)} \|u\|_d \|v\|_{t+q}.$$

Therefore, we conclude

$$\|u - u_J\|_{q-t} = \sup_{v \in H^{t+q}(\Gamma)} \frac{\langle A(u - u_J), v \rangle}{\|v\|_{t+q}} \lesssim 2^{J(q-d-t)} \|u\|_d,$$

which finishes the proof. \square

Note that in the extreme case $t = d - q$ we obtain the best possible convergence rate of the Galerkin scheme (3.6).

11. Complexity. In this section, we present a general theorem which shows that the overall complexity of assembling the compressed system matrix with sufficient accuracy can be kept of the order $\mathcal{O}(N_J)$, even when a computational cost of logarithmic order is allowed for each entry. This theorem is used in [19, 22] as the essential ingredient to provide a quadrature strategy which scales linearly.

THEOREM 11.1. *Assume that \mathbf{A}_J^ϵ is obtained by compressing the system matrix $\mathbf{A}_J = [\langle A\psi_{j',k'}, \psi_{j,k} \rangle]_{j_0 \leq j, j' < J, k \in \nabla_j, k' \in \nabla_{j'}}$ according to (7.1). The complexity of*

computing this compressed matrix is $\mathcal{O}(N_J)$ provided that for some $\alpha \geq 0$ at most $\mathcal{O}([J - \frac{j+j'}{2}]^\alpha)$ operations are spent on the approximate calculation of the nonvanishing entries $\langle A\psi_{j',k'}, \psi_{j,k} \rangle$.

Proof. (i) We begin with some technical preparations. Recall from the proof of Theorem 8.3 that the cut-off parameter of the first compression is given by

$$\mathcal{B}_{j,j'} \sim \max \{ 2^{-\min\{j,j'\}}, 2^{-J} 2^{(J-j)M} 2^{(J-j')M} \},$$

where, as in the proof of Theorem 8.3, $M = \frac{d'+\tilde{d}}{2(d+q)} < 1$. Moreover, we set $M' := \frac{2d'-2q}{d+d'}$ and $N' := \frac{\tilde{d}+d'}{d+2q}$ with d' given by (7.2). Note that M' and N' satisfy the relations $0 < M' < 1$ and $0 < N'$. As one readily verifies, the cut-off parameter with respect to the second compression may now be rewritten as

$$(11.1) \quad \mathcal{B}'_{j,j'} \sim \max \{ 2^{-j}, 2^{-j'} 2^{[JM'+(1-M')j'-j]N'} \}, \quad j \geq j'.$$

Further, we make use of the inequality $x^\alpha \lesssim 2^{2\delta x}$ which holds for all $x > 0$ and any fixed $\alpha, \delta > 0$. Thus, it suffices to prove the claim for $\mathcal{O}([J - \frac{j+j'}{2}]^\alpha)$ replaced by $\mathcal{O}(2^{\delta(J-j)} 2^{\delta(J-j')})$ where δ is chosen sufficiently small.

(ii) First, we determine now the complexity $\mathcal{C}^{(1)}$ of computing, within the above cost allowance, all matrix entries found in the block matrices $\mathbf{A}_{j,j'}^\epsilon = [\mathbf{A}_J^\epsilon]_{(j,\nabla_j),(j',\nabla_{j'})}$ with $\mathcal{B}_{j,j'} \sim 2^{-J} 2^{(J-j)M} 2^{(J-j')M}$. In such blocks, we have to process all coefficients $\langle A\psi_{j',k'}, \psi_{j,k} \rangle$ with

$$(11.2) \quad \text{dist}(\Omega_{j,k}, \Omega_{j',k'}) \lesssim \text{dist}_{j,j'}^{(1)} := 2^{-J} 2^{(J-j)M} 2^{(J-j')M}.$$

In each block, we find only $\mathcal{O}([2^{j'} \text{dist}_{j,j'}^{(1)}]^n)$ entries satisfying (11.2) per row, and hence a total of $\mathcal{O}([2^{j+j'} \text{dist}_{j,j'}^{(1)}]^n)$. Summing over all blocks yields

$$\begin{aligned} \mathcal{C}^{(1)} &\lesssim \sum_{j,j'=0}^J 2^{(j+j')n} 2^{-Jn} 2^{(J-j)(M+\delta)n} 2^{(J-j')(M+\delta)n} \\ &= 2^{Jn} \sum_{j,j'=0}^J 2^{(J-j)(M+\delta-1)n} 2^{(J-j')(M+\delta-1)n} \lesssim 2^{Jn}, \end{aligned}$$

provided that δ is chosen so as to ensure $M + \delta < 1$.

(iii) It remains to show that the complexity for computing the omitted blocks is likewise $\mathcal{O}(N_J)$. Without loss of generality, we assume $j \geq j'$ in the remainder of this proof, since the roles of j and j' can be reversed. Observing that, because of $0 < M' < 1$, one has $0 < JM' + (1 - M')j' \leq J$, we consider first the blocks $\mathbf{A}_{j,j'}^\epsilon$ with $(j, j') \in S$, where the index set S is given by

$$(11.3) \quad S := \{(j, j') : 0 \leq j' \leq J, JM' + (1 - M')j' \leq j \leq J\}.$$

In these blocks, we estimate the complexity $\mathcal{C}^{(2)}$ required for the approximate computation of the matrix entries $\langle A\psi_{j',k'}, \psi_{j,k} \rangle$ satisfying the relation

$$(11.4) \quad \text{dist}(\Omega'_{j,k}, \Omega_{j',k'}) \lesssim \text{dist}_{j,j'}^{(2)} := 2^{-j'} 2^{[JM'+(1-M')j'-j]N'},$$

where we refer to expression (11.1) for $\mathcal{B}'_{j,j'}$. Since $\text{dist}_{j,j'}^{(2)} \leq 2^{-j'}$ for all $(j, j') \in S$, in each block one finds only $\mathcal{O}([2^{jn} 2^{-j'(n-1)} \text{dist}_{j,j'}^{(2)}])$ nontrivial matrix entries per

column with (11.4), and thus a total of $\mathcal{O}([2^{jn}2^{j'} \text{dist}_{j,j'}^{(2)}])$. Therefore, noting that the set S is equivalent to $S = \{(j, j') : JM' \leq j \leq J, 0 \leq j' \leq \frac{j-JM'}{1-M'}\}$, the complexity is bounded by

$$\begin{aligned} \mathcal{C}^{(2)} &\lesssim \sum_{j=JM'}^J \sum_{j'=0}^{\frac{j-JM'}{1-M'}} 2^{jn}2^{[JM'+(1-M')j'-j]N'} 2^{\delta(J-j)}2^{\delta(J-j')} \\ &= \sum_{j=JM'}^J 2^{jn}2^{[JM'-j]N'} 2^{\delta(J-j)}2^{\delta J} \sum_{j'=0}^{\frac{j-JM'}{1-M'}} 2^{j'[(1-M')N'-\delta]} \\ &\lesssim 2^{\delta J \frac{2-M'}{1-M'}} \sum_{j=0}^J 2^{j(n-\delta \frac{2-M'}{1-M'})} \lesssim 2^{Jn}. \end{aligned}$$

$\mathcal{C}^{(2)}$ estimates the complexity for those blocks with $(j, j') \in S$ when $\mathcal{B}'_{j,j'} \sim \text{dist}_{j,j'}^{(2)}$. But according to (11.1), the cut-off parameter $\mathcal{B}'_{j,j'}$ is bounded from below by 2^{-j} . In the case of $\mathcal{B}'_{j,j'} \sim 2^{-j}$ we find $\mathcal{O}([2^{jn}2^{j'} \text{dist}_{j,j'}^{(3)}])$ matrix entries $\langle A\psi_{j',k'}, \psi_{j,k} \rangle$ with $\text{dist}(\Omega'_{j,k}, \Omega'_{j',k'}) \lesssim \text{dist}_{j,j'}^{(3)} := 2^{-j}$. Arguing analogously as above, summing over all blocks with $(j, j') \in S$, one obtains

$$\begin{aligned} \mathcal{C}^{(3)} &\lesssim \sum_{j=JM'}^J \sum_{j'=0}^{\frac{j-JM'}{1-M'}} 2^{j(n-1)}2^{j'}2^{\delta(J-j)}2^{\delta(J-j')} = \sum_{j=JM'}^J 2^{j(n-1)}2^{\delta(2J-j)} \sum_{j'=0}^{\frac{j-JM'}{1-M'}} 2^{j'(1-\delta)} \\ &\lesssim 2^{\delta J \frac{2-M'}{1-M'}} \sum_{j=0}^J 2^{j(n-\delta \frac{2-M'}{1-M'})} \lesssim 2^{Jn}. \end{aligned}$$

(iv) Finally, we consider the blocks $\mathbf{A}_{j,j'}^\epsilon$ with $j \geq j'$ and $(j, j') \notin S$. In view of step (ii), it suffices to consider all entries $\langle A\psi_{j',k'}, \psi_{j,k} \rangle$ which fulfill

$$(11.5) \quad \text{dist}(\Omega_{j,k}, \Omega_{j',k'}) \lesssim \text{dist}^{(4)} := 2^{-\min\{j',j\}} = 2^{-j'}.$$

Each block $\mathbf{A}_{j,j'}^\epsilon$ consists of only $\mathcal{O}([2^j2^{j'} \text{dist}^{(4)}]^n)$ entries with (11.5). Hence, according to (11.3), the complexity $\mathcal{C}^{(4)}$ for the computation of these entries is

$$\begin{aligned} \mathcal{C}^{(4)} &\lesssim \sum_{j'=0}^J \sum_{j=j'}^{JM'+(1-M')j'} 2^{jn}2^{\delta(J-j)}2^{\delta(J-j')} \lesssim \sum_{j'=0}^J 2^{2\delta(J-j')}2^{j'n} \sum_{j=0}^{(J-j')M'} 2^{j(n-\delta)} \\ &\lesssim 2^{Jn} \sum_{j'=0}^J 2^{(J-j')((M'-1)(n-\delta)+\delta)} \lesssim 2^{Jn}, \end{aligned}$$

since $(M' - 1)(n - \delta) + \delta < 0$. This completes this proof. \square

REFERENCES

[1] M. BEBENDORF AND S. RJASANOW, *Adaptive low-rank approximation of collocation matrices*, Computing, 70 (2003), pp. 1–24.
 [2] G. BEYLKIN, R. COIFMAN, AND V. ROKHLIN, *The fast wavelet transform and numerical algorithms*, Comm. Pure Appl. Math., 44 (1991), pp. 141–183.

- [3] C. CANUTO, A. TABACCO, AND K. URBAN, *The wavelet element method, part I: Construction and analysis*, Appl. Comput. Harmon. Anal., 6 (1999), pp. 1–52.
- [4] P.G. CIARLET, *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978.
- [5] A. COHEN, I. DAUBECHIES, AND J.-C. FEAUVEAU, *Biorthogonal bases of compactly supported wavelets*, Pure Appl. Math., 45 (1992), pp. 485–560.
- [6] A. COHEN AND R. MASSON, *Wavelet adaptive method for second order elliptic problems—boundary conditions and domain decomposition*, Numer. Math., 86 (2000), pp. 193–238.
- [7] M. COSTABEL, *Boundary integral operators on Lipschitz domains: Elementary results*, SIAM J. Math. Anal., 19 (1988), pp. 613–626.
- [8] W. DAHMEN, *Wavelet and multiscale methods for operator equations*, Acta Numer., 6 (1997), pp. 55–228.
- [9] W. DAHMEN AND A. KUNOTH, *Multilevel preconditioning*, Numer. Math., 63 (1992), pp. 315–344.
- [10] W. DAHMEN, A. KUNOTH, AND K. URBAN, *Biorthogonal spline-wavelets on the interval—stability and moment conditions*, Appl. Comp. Harmon. Anal., 6 (1999), pp. 259–302.
- [11] W. DAHMEN, S. PRÖSSDORF, AND R. SCHNEIDER, *Wavelet approximation methods for periodic pseudodifferential equations. Part II. Fast solution and matrix compression*, Adv. Comput. Math., 1 (1993), pp. 259–335.
- [12] W. DAHMEN, S. PRÖSSDORF, AND R. SCHNEIDER, *Multiscale methods for pseudodifferential equations on smooth manifolds*, in Wavelets: Theory, Algorithms, and Applications (Taormina, 1993), Wavelet Anal. Appl. 5, Academic Press, San Diego, CA, 1994, pp. 385–424.
- [13] W. DAHMEN AND R. SCHNEIDER, *Composite wavelet bases for operator equations*, Math. Comput., 68 (1999), pp. 1533–1567.
- [14] W. DAHMEN AND R. SCHNEIDER, *Wavelets on manifolds I. Construction and domain decomposition*, SIAM J. Math. Anal., 31 (1999), pp. 184–230.
- [15] W. DAHMEN AND R. STEVENSON, *Element-by-element construction of wavelets satisfying stability and moment conditions*, SIAM J. Numer. Anal., 37 (1999), pp. 319–352.
- [16] L. GREENGARD AND V. ROKHLIN, *A fast algorithm for particle simulation*, J. Comput. Phys., 73 (1987), pp. 325–348.
- [17] W. HACKBUSCH AND B.N. KHOROMSKIJ, *A sparse \mathcal{H} -matrix arithmetic. II: Application to multidimensional problems*, Computing, 64 (2000), pp. 21–47.
- [18] W. HACKBUSCH AND Z.P. NOWAK, *On the fast matrix multiplication in the boundary element method by panel clustering*, Numer. Math., 54 (1989), pp. 463–491.
- [19] H. HARBRECHT, *Wavelet Galerkin Schemes for the Boundary Element Method in Three Dimensions*, Ph.D. thesis, Technische Universität Chemnitz, Germany, 2001.
- [20] H. HARBRECHT, M. KONIK, AND R. SCHNEIDER, *Fully discrete wavelet Galerkin schemes*, Eng. Anal. Bound. Elem., 27 (2003), pp. 423–437.
- [21] H. HARBRECHT AND R. SCHNEIDER, *Biorthogonal wavelet bases for the boundary element method*, Math. Nachr., 167–188 (2004), pp. 269–270.
- [22] H. HARBRECHT AND R. SCHNEIDER, *Wavelet Galerkin schemes for boundary integral equations—implementation and quadrature*, SIAM J. Sci. Comput., to appear.
- [23] Y. MEYER, *Ondelettes et Opérateurs 2: Opérateur de Calderón-Zygmund*, Hermann, Paris, 1990.
- [24] T. VON PETERSDORFF, R. SCHNEIDER, AND C. SCHWAB, *Multiwavelets for second kind integral equations*, SIAM J. Numer. Anal., 34 (1997), pp. 2212–2227.
- [25] T. VON PETERSDORFF AND C. SCHWAB., *Fully discretized multiscale Galerkin BEM*, in Multiscale Wavelet Methods for PDEs, W. Dahmen, A. Kurdila, and P. Oswald, eds., Academic Press, San Diego, CA, 1997, pp. 287–346.
- [26] A. RATHSFELD, *A quadrature algorithm for wavelet Galerkin methods*, in Über Waveletalgorithmen für die Randelementmethode, Habilitation thesis, Technische Universität Chemnitz, Germany, 2001.
- [27] R. SCHNEIDER, *Multiskalen- und Wavelet-Matrixkompression: Analysisbasierte Methoden zur Lösung großer vollbesetzter Gleichungssysteme*, B.G. Teubner, Stuttgart, 1998.
- [28] E.M. STEIN, *Harmonic Analysis*, Princeton University Press, Princeton, NJ, 2002.
- [29] M.E. TAYLOR, *Pseudodifferential Operators*, Princeton University Press, Princeton, NJ, 1981.
- [30] S. GOREINOV, E. TYRTISCHNIKOV, AND Y. YEREMIN, *Matrix-free iterative solution strategies for large dense linear systems*, Numer. Linear Algebra Appl., 4 (1997), pp. 273–294.
- [31] W. WENDLAND, *Boundary element methods and their asymptotic convergence*, in Theoretical Acoustics and Numerical Techniques, CISM Courses and Lectures 277, P. Filippi, ed., Springer, New York, 1983, pp. 135–216.